arXiv:math/0511573v1 [math.NT] 23 Nov 2005

A NOTE ON THE FOURIER TRANSFORM OF p-ADIC q-INTEGRALS

Taekyun Kim

Department of Mathematics Education
Kongju University, Kongju 314-701, Korea
e-mail:tkim@kongju.ac.kr

Abstract. The p-adic q-integral ($= I_q$-integral) was defined by author in the previous paper [1, 3]. In this paper, we consider I_q-Fourier transform and investigate some properties which are related to this transform.

§1. Introduction

Let us denote $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{C}$ sets of positive integer, integer, rational and complex numbers respectively. Let p be prime and $x \in \mathbb{Q}$. Then $x = p^{v(x)}\frac{m}{n}$, where $m, n, v = v(x) \in \mathbb{Z}$, m and n are not divisible by p. Let $|x|_p = p^{-v(x)}$ and $|0|_p = 0$. Then $|x|_p$ is valuation on $\mathbb{Q}$ satisfying

$$|x+y|_p \leq \max\{|x|_p, |y|_p\}.$$

Completion of $\mathbb{Q}$ with respect to $|\cdot|_p$ is denoted by $\mathbb{Q}_p$ and called the field of p-adic rational numbers. $\mathbb{C}_p$ is the completion of algebraic closure of $\mathbb{Q}_p$ and $\mathbb{Z}_p = \{x \in \mathbb{Q}_p \mid |x|_p \leq 1\}$ is called the ring of p-adic rational integers(see 1, 2, 3, 4]). Let l be a fixed integer and let p be a fixed prime number. We set

$$\begin{aligned}
&X_l = \varprojlim_N (\mathbb{Z}/lp^N\mathbb{Z}), \text{ and } X_1 = \mathbb{Z}_p, \\
&X^* = \bigcup_{\substack{0<a<lp \\ (a,p)=1}} (a + lp\mathbb{Z}_p), \\
&a + lp^N\mathbb{Z}_p = \{x \in X \mid x \equiv a \pmod{lp^N}\},
\end{aligned}$$

where $N \in \mathbb{N}$ and $a \in \mathbb{Z}$ lies in $0 \leq a < lp^N$, cf. [3].

When one talks of q-extension, q is considered in many ways such as an indeterminate, a complex number $q \in \mathbb{C}$, or a p-adic number $q \in \mathbb{C}_p$. In this paper, we assume that $q \in \mathbb{C}_p$ with $|q-1|_p < p^{-\frac{1}{p-1}}$,

---

1991 *Mathematics Subject Classification.* 11B68, 11S80.

Key words and phrases. p-adic q-integrals, p-adic invariant integral on $\mathbb{Z}_p$, q-Volkenborn integral.

Typeset by $\mathcal{AMS}$-TEX

so that $q^x = \exp(x \log q)$ for each $x \in X$. We use the notation as $[x] = [x; q] = \frac{1-q^x}{1-q}$ for each $x \in X$. Hence $\lim_{q\to 1}[x] = x$, cf.[3]. For any positive integer N, we set

$$\mu_q(a + lp^N \mathbb{Z}_p) = \frac{q^a}{[lp^N]}, \text{ cf. [3]},$$

and this can be extended to a distribution on X. This distribution yields an integral as follows (see [3]):

$$I_q(f) = \int_{\mathbb{Z}_p} f(x)\, d\mu_q(x) = \int_X f(x)\, d\mu_q(x),$$

where $f \in UD(\mathbb{Z}_p) =$ the space of uniformly differentiable function on $\mathbb{Z}_p$ with values in $\mathbb{C}_p$, cf. [3]. Let C_{p^n} be the cyclic group consisting of all p^n-th roots of unity in $\mathbb{C}_p$ for any $n \geq 0$ and T_p be the direct limit of C_{p^n} with respect to the natural morphism, hence T_p ia the union of all C_{p^n} with discrete topology. U_p denotes the group of all principal units in $\mathbb{C}_p$. For any $f \in UD(\mathbb{Z}_p, \mathbb{C}_p)$, we have an integral $I_0(f)$ with respect to the so called invariant measure μ_0:

$$I_0(f) = \int_{\mathbb{Z}_p} f(x) d\mu_0(x) = \lim_{n\to\infty} \frac{1}{p^n} \sum_{x=0}^{p^n-1} f(x), \text{ cf. [3, 4]} ,$$

and the Fourier transform $\hat{f_w} = I_0(f\phi_w)$, where ϕ_w denotes a uniformly differentiable function on $\mathbb{Z}_p$ belonging to $w \in T_p$ defined by $\phi_w(x) = w^x$, cf. [4]. Now we introduce the convolution for any $f, g \in UD(\mathbb{Z}_p, \mathbb{C}_p)$ due to Woodcock as follows:

$$f * g(x) = \sum_w \hat{f_w} \hat{g_w} \phi_{w^{-1}}(x), \text{ see [3, 4]}.$$

As known results, $f * g \in UD(\mathbb{Z}_p, \mathbb{C}_p)$, and $\widehat{(f * g)}_w = \hat{f_w}\hat{g_w}$, (see [4]). In this paper, we consider I_q-Fourier transform and investigate some properties which are related to this transform. It is easy to see that I_q-Fourier transform is exactly same I_0-Fourier transform when $q = 1$.

§2. I_q-FOURIER TRANSFORM

For any $f \in UD(X)$, the p-adic q-integral was defined by

$$I_q(f) = \int_X f(x)\, d\mu_q(x) = \lim_{N\to\infty} \frac{1}{[lp^N]} \sum_{0 \leq x < lp^N} f(x) q^x, \text{ cf. [3]} .$$

Note that

$$I_0(f) = \lim_{q\to 1} I_q(f) = \int_{\mathbb{Z}_p} f(x)\, d\mu_1(x) = \lim_{N\to\infty} \frac{1}{lp^N} \sum_{0\leq x<lp^N} f(x),$$

and that

$$I_0(f_1) = I_0(f) + f'(0), \text{ where } f'(0) = \frac{d}{dx} f(x)|_{x=0} \text{ and } f_1(x) = f(x + 1).$$

Let $T_p = \cup_{n\geq 1} C_{p^n} = \lim_{n\to\infty} \mathbb{Z}/p^n\mathbb{Z}$, where $C_{p^n} = \{\xi \in X| \ \xi^{p^n} = 1\}$ is the cyclic group of order p^n, see [1]. For $\xi \in T_p$, we denote by $\phi_\xi : \mathbb{Z}_p \to \mathbb{C}_p$ the locally constant function $x \mapsto \xi^x$. If we take $f(x) = \phi_\xi(x)e^{tx}$, then we have that $\int_X e^{tx}\phi_\xi(x)d\mu_q(x) = \frac{t+\log q}{q\xi e^t - 1}$, cf. [2]. We now consider I_q-Fourier transform as follows:

$$(\hat{f_w})_q = I_q(\phi_w f) = \int_{\mathbb{Z}_p} \phi_w(x)f(x)d\mu_q(x), \text{ where } f \in UD(\mathbb{Z}_p),\ w \in T_p\ ,$$

and its inverse transform is derived by

$$\frac{\log q}{q-1}\lim_{n\to\infty}\sum_{x=0}^{p^n-1} w^{-x} I_q(\phi_w f) = \frac{\log q}{q-1}\lim_{n\to\infty}\sum_{x=0}^{p^n-1} w^{-x}\frac{1}{[p^n]}\sum_{z=0}^{p^n-1} w^z f(z)q^z = f(x)\phi_q(x).$$

Thus, we obtain the below proposition.

Proposition 1. *Let $f \in UD(\mathbb{Z}_p, \mathbb{C}_p)$. Then we have the inverse formula of I_q-Fourier transform as follows:*

$$f(x)\phi_q(x) = \frac{\log q}{q-1}\sum_w \phi_{w^{-1}}(\hat{f_w})_q, \ \textit{where} \sum_w = \lim_{n\to\infty}\sum_{w\in C_{p^n}}.$$

Remark. In [4], we note that if $\alpha \in U_p$, then ϕ_α is called locally analytic character and if $\alpha \in T_p$, then ϕ_α is called locally constant function. For $f, g \in UD(\mathbb{Z}_p)$, we consider the convolution of f, g by

$$f *_q g = \sum_w (\widehat{f_{wq^{-1}}})_q(\widehat{g_{wq^{-1}}})_q\phi_{w^{-1}}.$$

Thus, we note that

$$f *_q g \in UD(\mathbb{Z}_p), \text{ and } ((\widehat{f *_q g)_{wq^{-1}}})_q = (\widehat{f_{wq^{-1}}})_q(\widehat{g_{wq^{-1}}})_q.$$

And we also see that

$$UD(\mathbb{Z}_p, \mathbb{C}_p)/\{f \in UD(\mathbb{Z}_p, \mathbb{C}_p)| f' = 0\} \cong C(\mathbb{Z}_p, \mathbb{C}_p),$$

where $C(\mathbb{Z}_p, \mathbb{C}_p)$ is the space of the continuous function from $\mathbb{Z}_p$ to $\mathbb{C}_p$. Another convolution $\otimes_q$ is induced from the above convolution $*_q$ by $(f *_q g)' = -f' \otimes_q g'$. Then, we also see that $f \otimes_q g \in UD(\mathbb{Z}_p, \mathbb{C}_p)$. From these definitions, we can derive the below theorem.

Theorem 2. *For $f, g \in UD(\mathbb{Z}_p, \mathbb{C}_p)$, we have*

$$f *_q g(z) = \frac{q-1}{\log q} I_q^{(x)}(f(x)g(z-x)\phi_{q^{-1}}(x)) - (f \otimes_q g'(z)),$$

where $I_q^{(x)}$ means the integration with respect to the variable x.

Since $((\widehat{f *_q g)_{wq^{-1}}})_q = (\widehat{f_{wq^{-1}}})_q(\widehat{g_{wq^{-1}}})_q$, for $w \in T_p$, we have

$$\int_{\mathbb{Z}_p} (f *_q g(x))q^{-x}d\mu_q(x) = \int_{\mathbb{Z}_p} f(x)q^{-x}d\mu_q(x)\int_{\mathbb{Z}_p} g(x)q^{-x}d\mu_q(x).$$

From this, we can derive the below worthwhile and interesting formula:

Theorem 3. *Let $f, g \in UD(\mathbb{Z}_p, \mathbb{C}_p)$. Then we have*

$$I_q^{(z)}(f \otimes g'(z)q^{-z}) = \frac{q-1}{\log q} I_q^{(z)}(I_q^{(x)}(f(x)g(z-x)q^{-x})q^{-z}) - I_q^{(z)}(f(z)q^{-z}) I_q^{(z)}(g(z)q^{-z}).$$

REFERENCES

1. T. Kim, *On a q-analogue of the p-adic log-gamma functions and related integrals*, J. Number Theory **76** (1999), 320-329.
2. T. Kim, *A new approach to q-zeta function*, Advan. Stud. Contemp. Math. **11(2)** (2005), 157-162.
3. T.Kim, *q-Volkenborn Integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
4. C. F. Woodcock, *Convolutions on the ring of p-adic integers*, J. London Math. Soc. **20(2)** (1979), 101-108.